

Torus knot groups, Garside groups, complex reflection groups

Thomas Gobet

Institut Denis Poisson, Université de Tours

Conference *Braids and beyond*,
Université de Caen,
10th September 2021.


Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups



Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Question 30. Does the submonoid of B_n generated by $\sigma_1, \sigma_1\sigma_2, \dots, \sigma_1\sigma_2\cdots\sigma_{n-1}$ admit a finite presentation? Is it a Garside monoid?

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Question 30. Does the submonoid of B_n generated by $\sigma_1, \sigma_1\sigma_2, \dots, \sigma_1\sigma_2\cdots\sigma_{n-1}$ admit a finite presentation? Is it a Garside monoid?

Let Σ_n be the submonoid of the n -strand braid group \mathcal{B}_n generated by $\sigma_1, \sigma_1\sigma_2, \dots, \sigma_1\sigma_2\cdots\sigma_{n-1}$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Part I: Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

Part II: A new Garside structure on torus knot groups and some braid groups of complex reflection groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

Part III: "Reflection" quotients of torus knot groups

"Reflection" quotients of torus knot groups

Part I

Dehornoy-Digne-Godelle-Krammer- Michel's monoid and some torus knot groups

Garside structures: focusing on \mathcal{B}_3

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Garside structures: focusing on \mathcal{B}_3

- ▶ Four Garside structures on \mathcal{B}_3 :

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Garside structures: focusing on \mathcal{B}_3

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- Four Garside structures on \mathcal{B}_3 :

$$\left\langle \sigma_1, \sigma_2 \mid \begin{array}{l} \text{"Classical"} \\ \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \end{array} \right\rangle \quad \left\langle x, y \mid \begin{array}{l} \text{"Toric"} \\ x^2 = y^3 \end{array} \right\rangle$$

$$\left\langle \tau_1, \tau_2, \tau_3 \mid \begin{array}{l} \text{"Dual"} \\ \tau_1\tau_2 = \tau_2\tau_3 = \tau_3\tau_1 \end{array} \right\rangle \quad \left\langle a, b \mid \begin{array}{l} \text{"Exotic"} \\ aba = b^2 \end{array} \right\rangle$$

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

- ▶ Four Garside structures on \mathcal{B}_3 :

$$\left\langle \sigma_1, \sigma_2 \mid \begin{array}{l} \text{"Classical"} \\ \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \end{array} \right\rangle \quad \left\langle x, y \mid \begin{array}{l} \text{"Toric"} \\ x^2 = y^3 \end{array} \right\rangle$$

$$\left\langle \tau_1, \tau_2, \tau_3 \mid \begin{array}{l} \text{"Dual"} \\ \tau_1\tau_2 = \tau_2\tau_3 = \tau_3\tau_1 \end{array} \right\rangle \quad \left\langle a, b \mid \begin{array}{l} \text{"Exotic"} \\ aba = b^2 \end{array} \right\rangle$$

- ▶ **Dual braid monoid** : in terms of the classical Artin generators we have $\tau_1 = \sigma_1$, $\tau_2 = \sigma_2$, $\tau_3 = \sigma_1\sigma_2\sigma_1^{-1}$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

- ▶ Four Garside structures on \mathcal{B}_3 :

$$\left\langle \sigma_1, \sigma_2 \mid \begin{array}{l} \text{"Classical"} \\ \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \end{array} \right\rangle \quad \left\langle x, y \mid \begin{array}{l} \text{"Toric"} \\ x^2 = y^3 \end{array} \right\rangle$$

$$\left\langle \tau_1, \tau_2, \tau_3 \mid \begin{array}{l} \text{"Dual"} \\ \tau_1\tau_2 = \tau_2\tau_3 = \tau_3\tau_1 \end{array} \right\rangle \quad \left\langle a, b \mid \begin{array}{l} \text{"Exotic"} \\ aba = b^2 \end{array} \right\rangle$$

- ▶ **Dual braid monoid** : in terms of the classical Artin generators we have $\tau_1 = \sigma_1$, $\tau_2 = \sigma_2$, $\tau_3 = \sigma_1\sigma_2\sigma_1^{-1}$.
- ▶ **Toric monoid** : in terms of the classical Artin generators we have $x = \sigma_1\sigma_2\sigma_1$, $y = \sigma_1\sigma_2$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

- ▶ Four Garside structures on \mathcal{B}_3 :

$$\left\langle \sigma_1, \sigma_2 \mid \begin{array}{l} \text{"Classical"} \\ \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \end{array} \right\rangle \quad \left\langle x, y \mid \begin{array}{l} \text{"Toric"} \\ x^2 = y^3 \end{array} \right\rangle$$

$$\left\langle \tau_1, \tau_2, \tau_3 \mid \begin{array}{l} \text{"Dual"} \\ \tau_1\tau_2 = \tau_2\tau_3 = \tau_3\tau_1 \end{array} \right\rangle \quad \left\langle a, b \mid \begin{array}{l} \text{"Exotic"} \\ aba = b^2 \end{array} \right\rangle$$

- ▶ **Dual braid monoid** : in terms of the classical Artin generators we have $\tau_1 = \sigma_1$, $\tau_2 = \sigma_2$, $\tau_3 = \sigma_1\sigma_2\sigma_1^{-1}$.
- ▶ **Toric monoid** : in terms of the classical Artin generators we have $x = \sigma_1\sigma_2\sigma_1$, $y = \sigma_1\sigma_2$.
- ▶ **Exotic monoid** : in terms of the classical Artin generators we have $a = \sigma_1$, $b = \sigma_1\sigma_2$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

- ▶ Four Garside structures on \mathcal{B}_3 :

$$\begin{array}{cc} \begin{array}{c} \text{"Classical"} \\ \left\langle \sigma_1, \sigma_2 \mid \sigma_1 \sigma_2 \sigma_1 = \sigma_2 \sigma_1 \sigma_2 \right\rangle \end{array} & \begin{array}{c} \text{"Toric"} \\ \left\langle x, y \mid x^2 = y^3 \right\rangle \end{array} \\ \begin{array}{c} \text{"Dual"} \\ \left\langle \tau_1, \tau_2, \tau_3 \mid \tau_1 \tau_2 = \tau_2 \tau_3 = \tau_3 \tau_1 \right\rangle \end{array} & \begin{array}{c} \text{"Exotic"} \\ \left\langle a, b \mid aba = b^2 \right\rangle \end{array} \end{array}$$

- ▶ **Dual braid monoid** : in terms of the classical Artin generators we have $\tau_1 = \sigma_1$, $\tau_2 = \sigma_2$, $\tau_3 = \sigma_1 \sigma_2 \sigma_1^{-1}$.
- ▶ **Toric monoid** : in terms of the classical Artin generators we have $x = \sigma_1 \sigma_2 \sigma_1$, $y = \sigma_1 \sigma_2$.
- ▶ **Exotic monoid** : in terms of the classical Artin generators we have $a = \sigma_1$, $b = \sigma_1 \sigma_2$. The Garside element Δ is given by b^3 . We have $\text{Div}(\Delta) = \{1, a, b, ab, b^2, ba, bab, b^3\}$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Garside structures: focusing on \mathcal{B}_3

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Garside structures: focusing on \mathcal{B}_3

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The two Garside structures in the first column can be generalized to \mathcal{B}_n , $n > 3$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Garside structures: focusing on \mathcal{B}_3

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The two Garside structures in the first column can be generalized to \mathcal{B}_n , $n > 3$.
- ▶ The classical, dual, and toric structures can be generalized to *torus knot groups*.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

- ▶ The two Garside structures in the first column can be generalized to \mathcal{B}_n , $n > 3$.
- ▶ The classical, dual, and toric structures can be generalized to *torus knot groups*. Given n and m two coprime integers ≥ 2 , the group $G(n, m) = \langle x, y \mid x^n = y^m \rangle$ is the fundamental group of the complement of the torus knot $T_{n,m}$. It is a Garside group (Dehornoy-Paris 1999, Picantin 2003). One has $G(2, 3) \cong \mathcal{B}_3 \cong G(3, 2)$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

- ▶ The two Garside structures in the first column can be generalized to \mathcal{B}_n , $n > 3$.
- ▶ The classical, dual, and toric structures can be generalized to *torus knot groups*. Given n and m two coprime integers ≥ 2 , the group $G(n, m) = \langle x, y \mid x^n = y^m \rangle$ is the fundamental group of the complement of the torus knot $T_{n,m}$. It is a Garside group (Dehornoy-Paris 1999, Picantin 2003). One has $G(2, 3) \cong \mathcal{B}_3 \cong G(3, 2)$.
- ▶ What about the "exotic" structure ?

Dehornoy-Digne-Godolle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

► Let $n \geq 2$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ Let $n \geq 2$. Consider the monoid $\mathcal{M}(n)$ (resp. the group $\mathcal{G}(n)$) defined by the presentation

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ Let $n \geq 2$. Consider the monoid $\mathcal{M}(n)$ (resp. the group $\mathcal{G}(n)$) defined by the presentation

$$\left\langle \rho_1, \rho_2, \dots, \rho_n \mid \rho_1 \rho_n \rho_i = \rho_{i+1} \rho_n \text{ for all } 1 \leq i < n \right\rangle$$

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

- ▶ Let $n \geq 2$. Consider the monoid $\mathcal{M}(n)$ (resp. the group $\mathcal{G}(n)$) defined by the presentation

$$\left\langle \rho_1, \rho_2, \dots, \rho_n \mid \rho_1 \rho_n \rho_i = \rho_{i+1} \rho_n \text{ for all } 1 \leq i < n \right\rangle$$

- ▶ If $n = 2$, we obtain the presentation

$$\left\langle \rho_1, \rho_2 \mid \rho_1 \rho_2 \rho_1 = \rho_2^2 \right\rangle = \left\langle a, b \mid aba = b^2 \right\rangle$$

of the exotic monoid.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem

Let $n \geq 2$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem

Let $n \geq 2$.

1. *The monoid $\mathcal{M}(n)$ is a Garside monoid with (central) Garside element $\Delta = \rho_n^{n+1}$ and (left and right) lcm of the atoms equal to ρ_n^n .*

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem

Let $n \geq 2$.

1. The monoid $\mathcal{M}(n)$ is a Garside monoid with (central) Garside element $\Delta = \rho_n^{n+1}$ and (left and right) lcm of the atoms equal to ρ_n^n .
2. The Garside group $\mathcal{G}(n)$ is isomorphic to $G(n, n+1)$, the knot group of the torus knot $T_{n,n+1}$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem

Let $n \geq 2$.

1. The monoid $\mathcal{M}(n)$ is a Garside monoid with (central) Garside element $\Delta = \rho_n^{n+1}$ and (left and right) lcm of the atoms equal to ρ_n^n .
2. The Garside group $\mathcal{G}(n)$ is isomorphic to $G(n, n+1)$, the knot group of the torus knot $T_{n,n+1}$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

In fact, the group $\mathcal{G}(n) \cong G(n, n+1)$ surjects onto \mathcal{B}_{n+1} ($\rho_i \mapsto \sigma_1 \sigma_2 \cdots \sigma_i$). We have

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem

Let $n \geq 2$.

1. The monoid $\mathcal{M}(n)$ is a Garside monoid with (central) Garside element $\Delta = \rho_n^{n+1}$ and (left and right) lcm of the atoms equal to ρ_n^n .
2. The Garside group $\mathcal{G}(n)$ is isomorphic to $G(n, n+1)$, the knot group of the torus knot $T_{n, n+1}$.

Dehornoy-Digne-Godolle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

In fact, the group $\mathcal{G}(n) \cong G(n, n+1)$ surjects onto \mathcal{B}_{n+1} ($\rho_i \mapsto \sigma_1 \sigma_2 \cdots \sigma_i$). We have

Proposition

The image of $\mathcal{M}(n)$ inside \mathcal{B}_{n+1} is exactly the monoid Σ_{n+1} of Dehornoy-Digne-Godolle-Krammer-Michel.

Generalization of the exotic Garside structure

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem

Let $n \geq 2$.

1. The monoid $\mathcal{M}(n)$ is a Garside monoid with (central) Garside element $\Delta = \rho_n^{n+1}$ and (left and right) lcm of the atoms equal to ρ_n^n .
2. The Garside group $\mathcal{G}(n)$ is isomorphic to $G(n, n+1)$, the knot group of the torus knot $T_{n,n+1}$.

Dehornoy-Digne-Godolle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

In fact, the group $\mathcal{G}(n) \cong G(n, n+1)$ surjects onto \mathcal{B}_{n+1} ($\rho_i \mapsto \sigma_1 \sigma_2 \cdots \sigma_i$). We have

Proposition

The image of $\mathcal{M}(n)$ inside \mathcal{B}_{n+1} is exactly the monoid Σ_{n+1} of Dehornoy-Digne-Godolle-Krammer-Michel. In particular Σ_{n+1} is a quotient of $\mathcal{M}(n)$.

Example : $n = 3$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Example : $n = 3$

The group $\mathcal{G}(3)$ is isomorphic to the braid group of the exceptional complex reflection group G_{12} . The lattice of divisors of Δ (for left-divisibility) is given by :

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet


Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Example : $n = 3$

The group $\mathcal{G}(3)$ is isomorphic to the braid group of the exceptional complex reflection group G_{12} . The lattice of divisors of Δ (for left-divisibility) is given by :



About the lattice of simples

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

About the lattice of simples

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Theorem (Rognerud-G., 2021)

About the lattice of simples

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem (Rognerud-G., 2021)

Let $\Delta_n := \rho_n^{n+1}$ be the Garside element in $\mathcal{M}(n)$. We have

$$|\text{Div}(\Delta_n)| = F_{2(n+1)},$$

where F_i is the i -th entry of the Fibonacci sequence.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Theorem (Rognerud-G., 2021)

Let $\Delta_n := \rho_n^{n+1}$ be the Garside element in $\mathcal{M}(n)$. We have

$$|\text{Div}(\Delta_n)| = F_{2(n+1)},$$

where F_i is the i -th entry of the Fibonacci sequence. Hence the number of simples is given by 8, 21, 55, 144, 377, 987,

About the lattice of simples

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Theorem (Rognerud-G., 2021)

Let $\Delta_n := \rho_n^{n+1}$ be the Garside element in $\mathcal{M}(n)$. We have

$$|\text{Div}(\Delta_n)| = F_{2(n+1)},$$

where F_i is the i -th entry of the Fibonacci sequence. Hence the number of simples is given by 8, 21, 55, 144, 377, 987,

(There is a description of the lattice of simples in terms of *Schroeder trees*).

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Let us come back to DDGKM's question :

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Let us come back to DDGKM's question : "Is the monoid Σ_n a Garside monoid ?

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Let us come back to DDGKM's question : "Is the monoid Σ_n a Garside monoid ? Is it finitely presented ?"

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Let us come back to DDGKM's question : "Is the monoid Σ_n a Garside monoid ? Is it finitely presented ?"

Let \mathcal{H}_{n+1} (resp. \mathcal{H}_{n+1}^+) be the quotient of $\mathcal{G}(n)$ (resp. of $\mathcal{M}(n)$) defined by the presentation

$$\left\langle \rho_1, \rho_2, \dots, \rho_n \mid \rho_1 \rho_j \rho_{i-1} = \rho_i \rho_j, \forall 2 \leq i \leq j \leq n \right\rangle$$

Dehornoy-Digne-
Godolle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Let us come back to DDGKM's question : "Is the monoid Σ_n a Garside monoid ? Is it finitely presented ?"

Let \mathcal{H}_{n+1} (resp. \mathcal{H}_{n+1}^+) be the quotient of $\mathcal{G}(n)$ (resp. of $\mathcal{M}(n)$) defined by the presentation

$$\left\langle \rho_1, \rho_2, \dots, \rho_n \mid \rho_1 \rho_j \rho_{i-1} = \rho_i \rho_j, \forall 2 \leq i \leq j \leq n \right\rangle$$

Proposition

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Let us come back to DDGKM's question : "Is the monoid Σ_n a Garside monoid ? Is it finitely presented ?"

Let \mathcal{H}_{n+1} (resp. \mathcal{H}_{n+1}^+) be the quotient of $\mathcal{G}(n)$ (resp. of $\mathcal{M}(n)$) defined by the presentation

$$\left\langle \rho_1, \rho_2, \dots, \rho_n \mid \rho_1 \rho_j \rho_{i-1} = \rho_i \rho_j, \forall 2 \leq i \leq j \leq n \right\rangle$$

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Proposition

1. *There are surjections $\mathcal{M}(n) \twoheadrightarrow \mathcal{H}_{n+1}^+ \twoheadrightarrow \Sigma_{n+1}$,*

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Let us come back to DDGKM's question : "Is the monoid Σ_n a Garside monoid ? Is it finitely presented ?"

Let \mathcal{H}_{n+1} (resp. \mathcal{H}_{n+1}^+) be the quotient of $\mathcal{G}(n)$ (resp. of $\mathcal{M}(n)$) defined by the presentation

$$\left\langle \rho_1, \rho_2, \dots, \rho_n \mid \rho_1 \rho_j \rho_{i-1} = \rho_i \rho_j, \forall 2 \leq i \leq j \leq n \right\rangle$$

Dehornoy-Digne-Godolle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Proposition

1. *There are surjections $\mathcal{M}(n) \twoheadrightarrow \mathcal{H}_{n+1}^+ \twoheadrightarrow \Sigma_{n+1}$,*
2. *The monoid Σ_{n+1} is an Ore monoid with group of fractions isomorphic to \mathcal{B}_{n+1} . It is not a Garside monoid when $n > 2$.*

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Let us come back to DDGKM's question : "Is the monoid Σ_n a Garside monoid ? Is it finitely presented ?"

Let \mathcal{H}_{n+1} (resp. \mathcal{H}_{n+1}^+) be the quotient of $\mathcal{G}(n)$ (resp. of $\mathcal{M}(n)$) defined by the presentation

$$\left\langle \rho_1, \rho_2, \dots, \rho_n \mid \rho_1 \rho_j \rho_{i-1} = \rho_i \rho_j, \forall 2 \leq i \leq j \leq n \right\rangle$$

Dehornoy-Digne-Godolle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Proposition

1. *There are surjections $\mathcal{M}(n) \twoheadrightarrow \mathcal{H}_{n+1}^+ \twoheadrightarrow \Sigma_{n+1}$,*
2. *The monoid Σ_{n+1} is an Ore monoid with group of fractions isomorphic to \mathcal{B}_{n+1} . It is not a Garside monoid when $n > 2$.*
3. *We have $\mathcal{H}_{n+1} \cong \mathcal{B}_{n+1}$.*

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Moreover we conjecture :

Conjecture

The monoid \mathcal{H}_{n+1}^+ is cancellative.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Moreover we conjecture :

Conjecture

The monoid \mathcal{H}_{n+1}^+ is cancellative.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Lemma

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Moreover we conjecture :

Conjecture

The monoid \mathcal{H}_{n+1}^+ is cancellative.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Lemma

The following properties are equivalent:

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Moreover we conjecture :

Conjecture

The monoid \mathcal{H}_{n+1}^+ is cancellative.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Lemma

The following properties are equivalent:

1. *The monoid \mathcal{H}_{n+1}^+ is cancellative,*

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Moreover we conjecture :

Conjecture

The monoid \mathcal{H}_{n+1}^+ is cancellative.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Lemma

The following properties are equivalent:

- 1. The monoid \mathcal{H}_{n+1}^+ is cancellative,*
- 2. We have $\Sigma_{n+1} \cong \mathcal{H}_{n+1}^+$.*

"Reflection"
quotients of torus
knot groups

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Moreover we conjecture :

Conjecture

The monoid \mathcal{H}_{n+1}^+ is cancellative.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Lemma

The following properties are equivalent:

- 1. The monoid \mathcal{H}_{n+1}^+ is cancellative,*
- 2. We have $\Sigma_{n+1} \cong \mathcal{H}_{n+1}^+$.*

In particular, we can negatively answer the first part of DDGKM's question.

About the monoid Σ_n

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Moreover we conjecture :

Conjecture

The monoid \mathcal{H}_{n+1}^+ is cancellative.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Lemma

The following properties are equivalent:

- 1. The monoid \mathcal{H}_{n+1}^+ is cancellative,*
- 2. We have $\Sigma_{n+1} \cong \mathcal{H}_{n+1}^+$.*

"Reflection"
quotients of torus
knot groups

In particular, we can negatively answer the first part of DDGKM's question. If the above conjecture holds, then we can positively answer the second part of DDGKM's question.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Part II

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

Torus knot groups and braid groups of complex reflection groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups and braid groups of complex reflection groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem (Generalization to arbitrary torus knot groups)

Let $n, m \geq 2$ be two relatively prime integers. The Garside structure introduced above for $\mathcal{G}(n) \cong G(n, n+1)$ can be generalized to all torus knot groups $G(n, m)$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups and braid groups of complex reflection groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem (Generalization to arbitrary torus knot groups)

Let $n, m \geq 2$ be two relatively prime integers. The Garside structure introduced above for $\mathcal{G}(n) \cong G(n, n+1)$ can be generalized to all torus knot groups $G(n, m)$. More precisely, let $m = qn + r$ be the euclidean division of m by n ;

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups and braid groups of complex reflection groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem (Generalization to arbitrary torus knot groups)

Let $n, m \geq 2$ be two relatively prime integers. The Garside structure introduced above for $\mathcal{G}(n) \cong G(n, n+1)$ can be generalized to all torus knot groups $G(n, m)$. More precisely, let $m = qn + r$ be the euclidean division of m by n ; Define the monoid $\mathcal{M}(n, m)$ (resp. the group $\mathcal{G}(n, m)$) by the presentation

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Torus knot groups and braid groups of complex reflection groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem (Generalization to arbitrary torus knot groups)

Let $n, m \geq 2$ be two relatively prime integers. The Garside structure introduced above for $\mathcal{G}(n) \cong G(n, n+1)$ can be generalized to all torus knot groups $G(n, m)$. More precisely, let $m = qn + r$ be the euclidean division of m by n ; Define the monoid $\mathcal{M}(n, m)$ (resp. the group $\mathcal{G}(n, m)$) by the presentation

$$\left\langle \rho_1, \rho_2, \dots, \rho_n \mid \begin{array}{l} \rho_r \rho_n^q \rho_{i-r} = \rho_i \rho_n^q \text{ if } r < i \leq n, \\ \rho_r \rho_n^q \rho_{n+i-r} = \rho_i \rho_n^{q+1} \text{ if } 1 \leq i < r. \end{array} \right\rangle$$

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups and braid groups of complex reflection groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Theorem (Generalization to arbitrary torus knot groups)

Let $n, m \geq 2$ be two relatively prime integers. The Garside structure introduced above for $\mathcal{G}(n) \cong G(n, n+1)$ can be generalized to all torus knot groups $G(n, m)$. More precisely, let $m = qn + r$ be the euclidean division of m by n ; Define the monoid $\mathcal{M}(n, m)$ (resp. the group $\mathcal{G}(n, m)$) by the presentation

$$\left\langle \rho_1, \rho_2, \dots, \rho_n \mid \begin{array}{l} \rho_r \rho_n^q \rho_{i-r} = \rho_i \rho_n^q \text{ if } r < i \leq n, \\ \rho_r \rho_n^q \rho_{n+i-r} = \rho_i \rho_n^{q+1} \text{ if } 1 \leq i < r. \end{array} \right\rangle$$

(we have $\mathcal{M}(n) = \mathcal{M}(n, n+1)$).

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups and braid groups of complex reflection groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups and braid groups of complex reflection groups

- ▶ A few torus knot groups can be obtained as Artin-Tits groups of spherical type or more generally as braid groups of certain irreducible complex reflection groups of rank two, namely (in Coxeter or Shephard-Todd notation):

Torus knot groups, Garside groups, complex reflection groups

Thomas Gobet

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Torus knot groups and braid groups of complex reflection groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ A few torus knot groups can be obtained as Artin-Tits groups of spherical type or more generally as braid groups of certain irreducible complex reflection groups of rank two, namely (in Coxeter or Shephard-Todd notation):

$$G(2, 3) \cong \mathcal{B}_3 \cong \mathcal{B}(G_4) \cong \mathcal{B}(G_8) \cong \mathcal{B}(G_{16}),$$

$$G(2, 5) \cong \mathcal{B}(G_{20}),$$

$$G(2, m) \cong \mathcal{B}(I_2(m)), \quad m \text{ odd},$$

$$G(3, 4) \cong \mathcal{B}(G_{12}), \quad G(3, 5) \cong \mathcal{B}(G_{22}).$$

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups and braid groups of complex reflection groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ A few torus knot groups can be obtained as Artin-Tits groups of spherical type or more generally as braid groups of certain irreducible complex reflection groups of rank two, namely (in Coxeter or Shephard-Todd notation):

$$G(2, 3) \cong \mathcal{B}_3 \cong \mathcal{B}(G_4) \cong \mathcal{B}(G_8) \cong \mathcal{B}(G_{16}),$$

$$G(2, 5) \cong \mathcal{B}(G_{20}),$$

$$G(2, m) \cong \mathcal{B}(I_2(m)), \quad m \text{ odd},$$

$$G(3, 4) \cong \mathcal{B}(G_{12}), \quad G(3, 5) \cong \mathcal{B}(G_{22}).$$

- ▶ Moreover, a Garside structure analogous to the one introduced above for torus knot groups can be constructed for a few additional braid groups of complex reflection groups of rank two which are not isomorphic to torus knot groups.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Analogous Garside structures

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Analogous Garside structures

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following monoid yields a Garside monoid for G_{13} :

$$\left\langle \rho_1, \rho_2, \rho_3 \mid \begin{array}{l} \rho_1 \rho_3 \rho_2 = \rho_3^2 \\ \rho_2 \rho_3 \rho_1 = \rho_3^2 \end{array} \right\rangle.$$

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Analogous Garside structures

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following monoid yields a Garside monoid for G_{13} :

$$\left\langle \rho_1, \rho_2, \rho_3 \mid \begin{array}{l} \rho_1 \rho_3 \rho_2 = \rho_3^2 \\ \rho_2 \rho_3 \rho_1 = \rho_3^2 \end{array} \right\rangle.$$

- ▶ It was shown by Bannai that $\mathcal{B}(G_{13}) \cong \mathcal{B}(I_2(6))$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Analogous Garside structures

- ▶ The following monoid yields a Garside monoid for G_{13} :

$$\left\langle \rho_1, \rho_2, \rho_3 \mid \begin{array}{l} \rho_1 \rho_3 \rho_2 = \rho_3^2 \\ \rho_2 \rho_3 \rho_1 = \rho_3^2 \end{array} \right\rangle.$$

- ▶ It was shown by Bannai that $\mathcal{B}(G_{13}) \cong \mathcal{B}(I_2(6))$.
- ▶ Somewhat surprisingly, this monoid can be generalized to all Artin-Tits groups of even dihedral type:

Proposition

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Analogous Garside structures

- ▶ The following monoid yields a Garside monoid for G_{13} :

$$\left\langle \rho_1, \rho_2, \rho_3 \mid \begin{array}{l} \rho_1 \rho_3 \rho_2 = \rho_3^2 \\ \rho_2 \rho_3 \rho_1 = \rho_3^2 \end{array} \right\rangle.$$

- ▶ It was shown by Bannai that $\mathcal{B}(G_{13}) \cong \mathcal{B}(I_2(6))$.
- ▶ Somewhat surprisingly, this monoid can be generalized to all Artin-Tits groups of even dihedral type:

Proposition

Let $n \geq 1$.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Analogous Garside structures

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following monoid yields a Garside monoid for G_{13} :

$$\left\langle \rho_1, \rho_2, \rho_3 \mid \begin{array}{l} \rho_1 \rho_3 \rho_2 = \rho_3^2 \\ \rho_2 \rho_3 \rho_1 = \rho_3^2 \end{array} \right\rangle.$$

- ▶ It was shown by Bannai that $\mathcal{B}(G_{13}) \cong \mathcal{B}(I_2(6))$.
- ▶ Somewhat surprisingly, this monoid can be generalized to all Artin-Tits groups of even dihedral type:

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Proposition

Let $n \geq 1$. The monoid

$$\left\langle \tau_1, \tau_2, \rho \mid \begin{array}{l} \tau_1 \rho \tau_2 = \rho^2 \\ \tau_2 \rho^n \tau_1 = \rho^{n+1} \end{array} \right\rangle$$

is a Garside monoid for $\mathcal{B}(I_2(4 + 2n))$.

"Reflection"
quotients of torus
knot groups

Analogous Garside structures

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following monoid yields a Garside monoid for G_{13} :

$$\left\langle \rho_1, \rho_2, \rho_3 \mid \begin{array}{l} \rho_1 \rho_3 \rho_2 = \rho_3^2 \\ \rho_2 \rho_3 \rho_1 = \rho_3^2 \end{array} \right\rangle.$$

- ▶ It was shown by Bannai that $\mathcal{B}(G_{13}) \cong \mathcal{B}(I_2(6))$.
- ▶ Somewhat surprisingly, this monoid can be generalized to all Artin-Tits groups of even dihedral type:

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Proposition

Let $n \geq 1$. The monoid

$$\left\langle \tau_1, \tau_2, \rho \mid \begin{array}{l} \tau_1 \rho \tau_2 = \rho^2 \\ \tau_2 \rho^n \tau_1 = \rho^{n+1} \end{array} \right\rangle$$

is a Garside monoid for $\mathcal{B}(I_2(4 + 2n))$.

"Reflection"
quotients of torus
knot groups

- ▶ Note that the above monoid is still defined for $n = 0$. But we get $\rho = \tau_2 \tau_1$, and just recover the classical Garside structure on $\mathcal{B}(B_2)$.

Torus knot groups as "braid groups" of complex reflection groups ?

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups as "braid groups" of complex reflection groups ?

- ▶ Taking the quotient of $\mathcal{G}(2)$ by the relation $\rho_1^2 = 1$ (resp. $\rho_1^k = 1, k = 3, 4, 5$) yields the symmetric group \mathfrak{S}_3 (resp. the CRG G_4, G_8, G_{16}).

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups as "braid groups" of complex reflection groups ?

- ▶ Taking the quotient of $\mathcal{G}(2)$ by the relation $\rho_1^2 = 1$ (resp. $\rho_1^k = 1, k = 3, 4, 5$) yields the symmetric group \mathfrak{S}_3 (resp. the CRG G_4, G_8, G_{16}). Taking the quotient of $\mathcal{G}(3)$ by the relation $\rho_1^2 = 1$ yields the CRG G_{12} .

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups as "braid groups" of complex reflection groups ?

- ▶ Taking the quotient of $\mathcal{G}(2)$ by the relation $\rho_1^2 = 1$ (resp. $\rho_1^k = 1, k = 3, 4, 5$) yields the symmetric group \mathfrak{S}_3 (resp. the CRG G_4, G_8, G_{16}). Taking the quotient of $\mathcal{G}(3)$ by the relation $\rho_1^2 = 1$ yields the CRG G_{12} . Except for ρ_1 , the generators do not have a reflection as image.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups as "braid groups" of complex reflection groups ?

- ▶ Taking the quotient of $\mathcal{G}(2)$ by the relation $\rho_1^2 = 1$ (resp. $\rho_1^k = 1$, $k = 3, 4, 5$) yields the symmetric group \mathfrak{S}_3 (resp. the CRG G_4, G_8, G_{16}). Taking the quotient of $\mathcal{G}(3)$ by the relation $\rho_1^2 = 1$ yields the CRG G_{12} . Except for ρ_1 , the generators do not have a reflection as image. The quotient of $\mathcal{G}(4)$ by the relation $\rho_1^2 = 1$ is infinite.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups as "braid groups" of complex reflection groups ?

- ▶ Taking the quotient of $\mathcal{G}(2)$ by the relation $\rho_1^2 = 1$ (resp. $\rho_1^k = 1$, $k = 3, 4, 5$) yields the symmetric group \mathfrak{S}_3 (resp. the CRG G_4, G_8, G_{16}). Taking the quotient of $\mathcal{G}(3)$ by the relation $\rho_1^2 = 1$ yields the CRG G_{12} . Except for ρ_1 , the generators do not have a reflection as image. The quotient of $\mathcal{G}(4)$ by the relation $\rho_1^2 = 1$ is infinite.
- ▶ Let $n < m$ be as above.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups as "braid groups" of complex reflection groups ?

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ Taking the quotient of $\mathcal{G}(2)$ by the relation $\rho_1^2 = 1$ (resp. $\rho_1^k = 1$, $k = 3, 4, 5$) yields the symmetric group \mathfrak{S}_3 (resp. the CRG G_4, G_8, G_{16}). Taking the quotient of $\mathcal{G}(3)$ by the relation $\rho_1^2 = 1$ yields the CRG G_{12} . Except for ρ_1 , the generators do not have a reflection as image. The quotient of $\mathcal{G}(4)$ by the relation $\rho_1^2 = 1$ is infinite.
- ▶ Let $n < m$ be as above. The group $G(n, m)$ has a well-known presentation

$$\langle x_1, x_2, \dots, x_n \mid \underbrace{x_1 x_2 \cdots}_m = \underbrace{x_2 x_3 \cdots}_m = \cdots = \underbrace{x_n x_1 \cdots}_m \rangle.$$

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups as "braid groups" of complex reflection groups ?

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ Taking the quotient of $\mathcal{G}(2)$ by the relation $\rho_1^2 = 1$ (resp. $\rho_1^k = 1$, $k = 3, 4, 5$) yields the symmetric group \mathfrak{S}_3 (resp. the CRG G_4, G_8, G_{16}). Taking the quotient of $\mathcal{G}(3)$ by the relation $\rho_1^2 = 1$ yields the CRG G_{12} . Except for ρ_1 , the generators do not have a reflection as image. The quotient of $\mathcal{G}(4)$ by the relation $\rho_1^2 = 1$ is infinite.
- ▶ Let $n < m$ be as above. The group $G(n, m)$ has a well-known presentation

$$\langle x_1, x_2, \dots, x_n \mid \underbrace{x_1 x_2 \cdots}_m \text{ factors} = \underbrace{x_2 x_3 \cdots}_m \text{ factors} = \cdots = \underbrace{x_n x_1 \cdots}_m \text{ factors} \rangle.$$

- ▶ Taking reflection quotients as above corresponds to adding a relation $x_1^k = 1$ for $k \geq 2$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Torus knot groups as "braid groups" of complex reflection groups ?

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ Taking the quotient of $\mathcal{G}(2)$ by the relation $\rho_1^2 = 1$ (resp. $\rho_1^k = 1$, $k = 3, 4, 5$) yields the symmetric group \mathfrak{S}_3 (resp. the CRG G_4, G_8, G_{16}). Taking the quotient of $\mathcal{G}(3)$ by the relation $\rho_1^2 = 1$ yields the CRG G_{12} . Except for ρ_1 , the generators do not have a reflection as image. The quotient of $\mathcal{G}(4)$ by the relation $\rho_1^2 = 1$ is infinite.
- ▶ Let $n < m$ be as above. The group $G(n, m)$ has a well-known presentation

$$\langle x_1, x_2, \dots, x_n \mid \underbrace{x_1 x_2 \cdots}_m \text{ factors} = \underbrace{x_2 x_3 \cdots}_m \text{ factors} = \cdots = \underbrace{x_n x_1 \cdots}_m \text{ factors} \rangle.$$

- ▶ Taking reflection quotients as above corresponds to adding a relation $x_1^k = 1$ for $k \geq 2$. One then has $x_i^k = 1$ for all i .

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Torus knot groups as "braid groups" of complex reflection groups ?

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ Taking the quotient of $\mathcal{G}(2)$ by the relation $\rho_1^2 = 1$ (resp. $\rho_1^k = 1$, $k = 3, 4, 5$) yields the symmetric group \mathfrak{S}_3 (resp. the CRG G_4, G_8, G_{16}). Taking the quotient of $\mathcal{G}(3)$ by the relation $\rho_1^2 = 1$ yields the CRG G_{12} . Except for ρ_1 , the generators do not have a reflection as image. The quotient of $\mathcal{G}(4)$ by the relation $\rho_1^2 = 1$ is infinite.
- ▶ Let $n < m$ be as above. The group $G(n, m)$ has a well-known presentation

$$\langle x_1, x_2, \dots, x_n \mid \underbrace{x_1 x_2 \cdots}_m = \underbrace{x_2 x_3 \cdots}_m = \cdots = \underbrace{x_n x_1 \cdots}_m \rangle.$$

- ▶ Taking reflection quotients as above corresponds to adding a relation $x_1^k = 1$ for $k \geq 2$. One then has $x_i^k = 1$ for all i . Denote by $W(k, n, m)$ such a quotient.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Part III

"Reflection" quotients of torus knot groups

Particular cases of $W(k, n, m)$ and questions

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Particular cases of $W(k, n, m)$ and questions

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following groups appear as particular instances of the groups $W(k, n, m)$:

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Particular cases of $W(k, n, m)$ and questions

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following groups appear as particular instances of the groups $W(k, n, m)$:
 - ▶ $W(2, 2, m)$ is the dihedral group $I_2(m)$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Particular cases of $W(k, n, m)$ and questions

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following groups appear as particular instances of the groups $W(k, n, m)$:
 - ▶ $W(2, 2, m)$ is the dihedral group $I_2(m)$.
 - ▶ $W(2, 3, 4) = G_{12}$, $W(2, 3, 5) = G_{22}$, $W(3, 2, 5) = G_{20}$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Particular cases of $W(k, n, m)$ and questions

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following groups appear as particular instances of the groups $W(k, n, m)$:
 - ▶ $W(2, 2, m)$ is the dihedral group $I_2(m)$.
 - ▶ $W(2, 3, 4) = G_{12}$, $W(2, 3, 5) = G_{22}$, $W(3, 2, 5) = G_{20}$.
 - ▶ $W(k, 2, 3)$ ($k \geq 3$) is Coxeter's truncated braid group on two generators, i.e., the quotient of \mathcal{B}_3 by the relation $\sigma_1^k = \sigma_2^k = 1$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Particular cases of $W(k, n, m)$ and questions

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following groups appear as particular instances of the groups $W(k, n, m)$:
 - ▶ $W(2, 2, m)$ is the dihedral group $I_2(m)$.
 - ▶ $W(2, 3, 4) = G_{12}$, $W(2, 3, 5) = G_{22}$, $W(3, 2, 5) = G_{20}$.
 - ▶ $W(k, 2, 3)$ ($k \geq 3$) is Coxeter's truncated braid group on two generators, i.e., the quotient of \mathcal{B}_3 by the relation $\sigma_1^k = \sigma_2^k = 1$. We have $W(3, 2, 3) = G_4$, $W(4, 2, 3) = G_8$, $W(5, 2, 3) = G_{16}$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Particular cases of $W(k, n, m)$ and questions

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following groups appear as particular instances of the groups $W(k, n, m)$:
 - ▶ $W(2, 2, m)$ is the dihedral group $I_2(m)$.
 - ▶ $W(2, 3, 4) = G_{12}$, $W(2, 3, 5) = G_{22}$, $W(3, 2, 5) = G_{20}$.
 - ▶ $W(k, 2, 3)$ ($k \geq 3$) is Coxeter's truncated braid group on two generators, i.e., the quotient of \mathcal{B}_3 by the relation $\sigma_1^k = \sigma_2^k = 1$. We have $W(3, 2, 3) = G_4$, $W(4, 2, 3) = G_8$, $W(5, 2, 3) = G_{16}$. When $k \geq 6$ the group is infinite, and admits a representation as a complex reflection group on \mathbb{C}^2 (which is unfaithful in general).

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Particular cases of $W(k, n, m)$ and questions

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The following groups appear as particular instances of the groups $W(k, n, m)$:
 - ▶ $W(2, 2, m)$ is the dihedral group $I_2(m)$.
 - ▶ $W(2, 3, 4) = G_{12}$, $W(2, 3, 5) = G_{22}$, $W(3, 2, 5) = G_{20}$.
 - ▶ $W(k, 2, 3)$ ($k \geq 3$) is Coxeter's truncated braid group on two generators, i.e., the quotient of \mathcal{B}_3 by the relation $\sigma_1^k = \sigma_2^k = 1$. We have $W(3, 2, 3) = G_4$, $W(4, 2, 3) = G_8$, $W(5, 2, 3) = G_{16}$. When $k \geq 6$ the group is infinite, and admits a representation as a complex reflection group on \mathbb{C}^2 (which is unfaithful in general).
- ▶ Every finite reflection group occurring above is a rank-two complex reflection group.

Dehornoy-Digne-Godolle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

J -groups (Achar-Aubert, 2008)

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

J -groups (Achar-Aubert, 2008)

Achar and Aubert introduced a family of (in general infinite) groups, called J -groups.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

J -groups (Achar-Aubert, 2008)

Achar and Aubert introduced a family of (in general infinite) groups, called J -groups.

Let a, b, c three integers ≥ 1 .

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

J -groups (Achar-Aubert, 2008)

Achar and Aubert introduced a family of (in general infinite) groups, called J -groups.

Let a, b, c three integers ≥ 1 . Let $J \begin{pmatrix} a & b & c \end{pmatrix}$ be the group defined by the presentation

$$\langle s, t, u \mid s^a = t^b = u^c = 1, stu = tus = ust \rangle$$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

J -groups (Achar-Aubert, 2008)

Achar and Aubert introduced a family of (in general infinite) groups, called J -groups.

Let a, b, c three integers ≥ 1 . Let $J \begin{pmatrix} a & b & c \end{pmatrix}$ be the group defined by the presentation

$$\langle s, t, u \mid s^a = t^b = u^c = 1, stu = tus = ust \rangle$$

Let a', b' and c' be three pairwise coprime integers, dividing a, b and c respectively.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

J -groups (Achar-Aubert, 2008)

Achar and Aubert introduced a family of (in general infinite) groups, called J -groups.

Let a, b, c three integers ≥ 1 . Let $J \begin{pmatrix} a & b & c \end{pmatrix}$ be the group defined by the presentation

$$\langle s, t, u \mid s^a = t^b = u^c = 1, stu = tus = ust \rangle$$

Let a', b' and c' be three pairwise coprime integers, dividing a, b and c respectively. Let $J \begin{pmatrix} a & b & c \\ a' & b' & c' \end{pmatrix}$ be the normal subgroup of $J \begin{pmatrix} a & b & c \end{pmatrix}$ generated by $s^{a'}, t^{b'}$ and $u^{c'}$.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

J -groups (Achar-Aubert, 2008)

Achar and Aubert introduced a family of (in general infinite) groups, called J -groups.

Let a, b, c three integers ≥ 1 . Let $J \begin{pmatrix} a & b & c \\ & & \end{pmatrix}$ be the group defined by the presentation

$$\langle s, t, u \mid s^a = t^b = u^c = 1, stu = tus = ust \rangle$$

Let a', b' and c' be three pairwise coprime integers, dividing a, b and c respectively. Let $J \begin{pmatrix} a & b & c \\ a' & b' & c' \end{pmatrix}$ be the normal subgroup of $J \begin{pmatrix} a & b & c \\ & & \end{pmatrix}$ generated by $s^{a'}, t^{b'}$ and $u^{c'}$. We omit 1's in the second row of parameters, consistently with

$$J \begin{pmatrix} a & b & c \\ 1 & 1 & 1 \end{pmatrix} = J \begin{pmatrix} a & b & c \\ & & \end{pmatrix}.$$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

J -groups

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Theorem (Achar-Aubert, 2008)

A J -group is finite if and only if it is a finite complex reflection group of rank 2.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Theorem (Achar-Aubert, 2008)

A J -group is finite if and only if it is a finite complex reflection group of rank 2.

- ▶ Achar and Aubert also showed that every J -group G admits a representation $\rho : G \rightarrow \mathrm{GL}_2(\mathbb{C})$, where $\rho(s)$, $\rho(t)$ and $\rho(u)$ are reflections preserving a Hermitian form (so that the image is a complex reflection group).

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Theorem (Achar-Aubert, 2008)

A J -group is finite if and only if it is a finite complex reflection group of rank 2.

- ▶ Achar and Aubert also showed that every J -group G admits a representation $\rho : G \rightarrow \mathrm{GL}_2(\mathbb{C})$, where $\rho(s)$, $\rho(t)$ and $\rho(u)$ are reflections preserving a Hermitian form (so that the image is a complex reflection group). When G is not finite, this representation is *not* faithful in general.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Theorem (Achar-Aubert, 2008)

A J -group is finite if and only if it is a finite complex reflection group of rank 2.

- ▶ Achar and Aubert also showed that every J -group G admits a representation $\rho : G \rightarrow \mathrm{GL}_2(\mathbb{C})$, where $\rho(s)$, $\rho(t)$ and $\rho(u)$ are reflections preserving a Hermitian form (so that the image is a complex reflection group). When G is not finite, this representation is *not* faithful in general.
- ▶ This result is reminiscent of the following theorem : let W be a Coxeter group. Then W is a real reflection group if and only if W is finite.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

A family of J -groups with a single conjugacy class of "reflecting hyperplanes"

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

A family of J -groups with a single conjugacy class of "reflecting hyperplanes"

Definition

Let k, n, m be three integers ≥ 2 with $n < m$ and n, m coprime.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

A family of J -groups with a single conjugacy class of "reflecting hyperplanes"

Definition

Let k, n, m be three integers ≥ 2 with $n < m$ and n, m coprime. Let $\mathcal{J}(n, m, k) := J \begin{pmatrix} k & n & m \\ & n & m \end{pmatrix}$.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

A family of J -groups with a single conjugacy class of "reflecting hyperplanes"

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Definition

Let k, n, m be three integers ≥ 2 with $n < m$ and n, m coprime. Let $\mathcal{J}(n, m, k) := J \begin{pmatrix} k & n & m \\ & n & m \end{pmatrix}$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Theorem

Let k, n, m as above. We have $W(k, n, m) \cong \mathcal{J}(k, n, m)$.

"Reflection"
quotients of torus
knot groups

A family of J -groups with a single conjugacy class of "reflecting hyperplanes"

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Definition

Let k, n, m be three integers ≥ 2 with $n < m$ and n, m coprime. Let $\mathcal{J}(n, m, k) := J \begin{pmatrix} k & n & m \\ & n & m \end{pmatrix}$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Theorem

Let k, n, m as above. We have $W(k, n, m) \cong \mathcal{J}(k, n, m)$.

- ▶ The group in the LHS is a quotient of the torus knot group $G(n, m)$.

"Reflection"
quotients of torus
knot groups

A family of J -groups with a single conjugacy class of "reflecting hyperplanes"

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Definition

Let k, n, m be three integers ≥ 2 with $n < m$ and n, m coprime. Let $\mathcal{J}(n, m, k) := J \begin{pmatrix} k & n & m \\ & n & m \end{pmatrix}$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Theorem

Let k, n, m as above. We have $W(k, n, m) \cong \mathcal{J}(k, n, m)$.

- ▶ The group in the LHS is a quotient of the torus knot group $G(n, m)$. When the J -group is finite, this group $G(n, m)$ is the braid group of the corresponding finite CRG.

"Reflection"
quotients of torus
knot groups

A family of J -groups with a single conjugacy class of "reflecting hyperplanes"

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Definition

Let k, n, m be three integers ≥ 2 with $n < m$ and n, m coprime. Let $\mathcal{J}(n, m, k) := J \begin{pmatrix} k & n & m \\ & n & m \end{pmatrix}$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

Theorem

Let k, n, m as above. We have $W(k, n, m) \cong \mathcal{J}(k, n, m)$.

"Reflection"
quotients of torus
knot groups

- ▶ The group in the LHS is a quotient of the torus knot group $G(n, m)$. When the J -group is finite, this group $G(n, m)$ is the braid group of the corresponding finite CRG.
- ▶ In this way, we associate a Garside group to every J -group in the above defined family, in such a way that whenever the J -group is finite, one recovers the braid group of the CRG.

Braid groups of J -groups ?

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Braid groups of J -groups ?

- ▶ From the above observations, it is tempting to call $G(n, m)$ the "braid group" of $W(k, n, m)$.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Braid groups of J -groups ?

- ▶ From the above observations, it is tempting to call $G(n, m)$ the "braid group" of $W(k, n, m)$.
- ▶ However, it is not clear that this combinatorial definition is the right one. Note that

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Braid groups of J -groups ?

- ▶ From the above observations, it is tempting to call $G(n, m)$ the "braid group" of $W(k, n, m)$.
- ▶ However, it is not clear that this combinatorial definition is the right one. Note that
 - ▶ When $W(k, n, m)$ is infinite, we do not know if these groups have a faithful representation as complex reflection groups of rank two in general.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Braid groups of J -groups ?

- ▶ From the above observations, it is tempting to call $G(n, m)$ the "braid group" of $W(k, n, m)$.
- ▶ However, it is not clear that this combinatorial definition is the right one. Note that
 - ▶ When $W(k, n, m)$ is infinite, we do not know if these groups have a faithful representation as complex reflection groups of rank two in general. Hence it is hard to define the "braid group" as the fundamental group of the space of regular orbits of a reflection representation.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Braid groups of J -groups ?

- ▶ From the above observations, it is tempting to call $G(n, m)$ the "braid group" of $W(k, n, m)$.
- ▶ However, it is not clear that this combinatorial definition is the right one. Note that
 - ▶ When $W(k, n, m)$ is infinite, we do not know if these groups have a faithful representation as complex reflection groups of rank two in general. Hence it is hard to define the "braid group" as the fundamental group of the space of regular orbits of a reflection representation.
 - ▶ For $(n_1, m_1) \neq (n_2, m_2)$ (with $n_i < m_i$ and n_i, m_i coprime), we have $G(n_1, m_1) \not\cong G(n_2, m_2)$ (Schreier, 1924).

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Braid groups of J -groups ?

- ▶ From the above observations, it is tempting to call $G(n, m)$ the "braid group" of $W(k, n, m)$.
- ▶ However, it is not clear that this combinatorial definition is the right one. Note that
 - ▶ When $W(k, n, m)$ is infinite, we do not know if these groups have a faithful representation as complex reflection groups of rank two in general. Hence it is hard to define the "braid group" as the fundamental group of the space of regular orbits of a reflection representation.
 - ▶ For $(n_1, m_1) \neq (n_2, m_2)$ (with $n_i < m_i$ and n_i, m_i coprime), we have $G(n_1, m_1) \not\cong G(n_2, m_2)$ (Schreier, 1924). It is not clear a priori that if $W(k_1, n_1, m_1) \cong W(k_2, n_2, m_2)$ (as "reflection groups"), then $(k_1, n_1, m_1) = (k_2, n_2, m_2)$.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Braid groups of J -groups ?

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ From the above observations, it is tempting to call $G(n, m)$ the "braid group" of $W(k, n, m)$.
- ▶ However, it is not clear that this combinatorial definition is the right one. Note that
 - ▶ When $W(k, n, m)$ is infinite, we do not know if these groups have a faithful representation as complex reflection groups of rank two in general. Hence it is hard to define the "braid group" as the fundamental group of the space of regular orbits of a reflection representation.
 - ▶ For $(n_1, m_1) \neq (n_2, m_2)$ (with $n_i < m_i$ and n_i, m_i coprime), we have $G(n_1, m_1) \not\cong G(n_2, m_2)$ (Schreier, 1924). It is not clear a priori that if $W(k_1, n_1, m_1) \cong W(k_2, n_2, m_2)$ (as "reflection groups"), then $(k_1, n_1, m_1) = (k_2, n_2, m_2)$.
- ▶ In fact the parameter k_i is the number of conjugacy classes of reflections in $W(k_i, n_i, m_i)$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Braid groups of J -groups ?

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ From the above observations, it is tempting to call $G(n, m)$ the "braid group" of $W(k, n, m)$.
- ▶ However, it is not clear that this combinatorial definition is the right one. Note that
 - ▶ When $W(k, n, m)$ is infinite, we do not know if these groups have a faithful representation as complex reflection groups of rank two in general. Hence it is hard to define the "braid group" as the fundamental group of the space of regular orbits of a reflection representation.
 - ▶ For $(n_1, m_1) \neq (n_2, m_2)$ (with $n_i < m_i$ and n_i, m_i coprime), we have $G(n_1, m_1) \not\cong G(n_2, m_2)$ (Schreier, 1924). It is not clear a priori that if $W(k_1, n_1, m_1) \cong W(k_2, n_2, m_2)$ (as "reflection groups"), then $(k_1, n_1, m_1) = (k_2, n_2, m_2)$.
- ▶ In fact the parameter k_i is the number of conjugacy classes of reflections in $W(k_i, n_i, m_i)$. Hence at least $W(k_1, n_1, m_1) \cong_{\text{ref}} W(k_2, n_2, m_2) \Rightarrow k_1 = k_2$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The element $c := (x_1 x_2 \cdots x_n)^m$ is central in $W(k, n, m)$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

- ▶ The element $c := (x_1 x_2 \cdots x_n)^m$ is central in $W(k, n, m)$. Let $\overline{W}(k, n, m) = W(k, n, m)/(c = 1)$.

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ The element $c := (x_1 x_2 \cdots x_n)^m$ is central in $W(k, n, m)$. Let $\overline{W}(k, n, m) = W(k, n, m)/(c = 1)$. Let $W_{k,n,m}$ be the rank three Coxeter group

$$W_{k,n,m} = \left\langle r_1, r_2, r_3 \mid \begin{array}{l} r_1^2 = r_2^2 = r_3^2 = 1, \\ (r_1 r_2)^k = (r_2 r_3)^n = (r_3 r_1)^m = 1 \end{array} \right\rangle$$

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

- ▶ The element $c := (x_1 x_2 \cdots x_n)^m$ is central in $W(k, n, m)$. Let $\overline{W}(k, n, m) = W(k, n, m)/(c = 1)$. Let $W_{k,n,m}$ be the rank three Coxeter group

$$W_{k,n,m} = \left\langle r_1, r_2, r_3 \mid \begin{array}{l} r_1^2 = r_2^2 = r_3^2 = 1, \\ (r_1 r_2)^k = (r_2 r_3)^n = (r_3 r_1)^m = 1 \end{array} \right\rangle$$

- ▶ There is a group morphism

$$J := J \begin{pmatrix} k & n & m \end{pmatrix} \longrightarrow W_{k,n,m}, \quad s \mapsto r_1 r_2, \quad t \mapsto r_2 r_3, \\ u \mapsto r_3 r_1.$$

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

- The element $c := (x_1 x_2 \cdots x_n)^m$ is central in $W(k, n, m)$. Let $\overline{W}(k, n, m) = W(k, n, m)/(c = 1)$. Let $W_{k,n,m}$ be the rank three Coxeter group

$$W_{k,n,m} = \left\langle r_1, r_2, r_3 \mid \begin{array}{l} r_1^2 = r_2^2 = r_3^2 = 1, \\ (r_1 r_2)^k = (r_2 r_3)^n = (r_3 r_1)^m = 1 \end{array} \right\rangle$$

- There is a group morphism

$$J := J \begin{pmatrix} k & n & m \end{pmatrix} \longrightarrow W_{k,n,m}, \quad s \mapsto r_1 r_2, \quad t \mapsto r_2 r_3,$$

$u \mapsto r_3 r_1$. Its image is the alternating subgroup $W_{k,n,m}^+$ of $W_{k,n,m}$. One has $J/(stu = 1) \cong W_{k,n,m}^+$.

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Proposition

We have $\overline{W}(k, n, m) \cong W_{k,n,m}^+$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Proposition

We have $\overline{W}(k, n, m) \cong W_{k,n,m}^+$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

Proposition

Let (W, S) be a Coxeter system of rank ≥ 3 . Then the center of the alternating subgroup W^+ of W is included in the center of W .

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Proposition

We have $\overline{W}(k, n, m) \cong W_{k,n,m}^+$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

Proposition

Let (W, S) be a Coxeter system of rank ≥ 3 . Then the center of the alternating subgroup W^+ of W is included in the center of W . In particular, if W is infinite irreducible with $|S| \geq 3$, then the center of W^+ is trivial.

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Proposition

We have $\overline{W}(k, n, m) \cong W_{k,n,m}^+$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

Proposition

Let (W, S) be a Coxeter system of rank ≥ 3 . Then the center of the alternating subgroup W^+ of W is included in the center of W . In particular, if W is infinite irreducible with $|S| \geq 3$, then the center of W^+ is trivial.

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Corollary

The center of $W(k, n, m)$ is cyclic, generated by c .

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Proposition

We have $\overline{W}(k, n, m) \cong W_{k,n,m}^+$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

Proposition

Let (W, S) be a Coxeter system of rank ≥ 3 . Then the center of the alternating subgroup W^+ of W is included in the center of W . In particular, if W is infinite irreducible with $|S| \geq 3$, then the center of W^+ is trivial.

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Corollary

The center of $W(k, n, m)$ is cyclic, generated by c .

- In general we do not know if c has finite order or not.

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

- ▶ In the cases where $W(k, n, m)$ is finite we recover the known description of $\overline{W}(k, n, m)$ as a particular case:

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ In the cases where $W(k, n, m)$ is finite we recover the known description of $\overline{W}(k, n, m)$ as a particular case:

| k | n | m | $W(k, n, m)$ | $\overline{W}(k, n, m)$ |
|-----|-----|-----|--------------|---------------------------------|
| 2 | 3 | 4 | G_{12} | $W(B_3)^+ \cong \mathfrak{S}_4$ |
| 2 | 3 | 5 | G_{22} | $W(H_3)^+ \cong \mathfrak{A}_5$ |
| 3 | 2 | 3 | G_4 | $W(A_3)^+ \cong \mathfrak{A}_3$ |
| 4 | 2 | 3 | G_8 | $W(B_3)^+ \cong \mathfrak{S}_4$ |
| 5 | 2 | 3 | G_{16} | $W(H_3)^+ \cong \mathfrak{A}_5$ |
| 3 | 2 | 5 | G_{20} | $W(H_3)^+ \cong \mathfrak{A}_5$ |

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Center of $W(k, n, m)$

Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

- ▶ In the cases where $W(k, n, m)$ is finite we recover the known description of $\overline{W}(k, n, m)$ as a particular case:

| k | n | m | $W(k, n, m)$ | $\overline{W}(k, n, m)$ |
|-----|-----|-----|--------------|---------------------------------|
| 2 | 3 | 4 | G_{12} | $W(B_3)^+ \cong \mathfrak{S}_4$ |
| 2 | 3 | 5 | G_{22} | $W(H_3)^+ \cong \mathfrak{A}_5$ |
| 3 | 2 | 3 | G_4 | $W(A_3)^+ \cong \mathfrak{A}_3$ |
| 4 | 2 | 3 | G_8 | $W(B_3)^+ \cong \mathfrak{S}_4$ |
| 5 | 2 | 3 | G_{16} | $W(H_3)^+ \cong \mathfrak{A}_5$ |
| 3 | 2 | 5 | G_{20} | $W(H_3)^+ \cong \mathfrak{A}_5$ |

Dehornoy-Digne-Godelle-Krammer-Michel's monoid and some torus knot groups

A new Garside structure on torus knot groups and some braid groups of complex reflection groups

"Reflection" quotients of torus knot groups

- ▶ With the observations above, to show that $W(k, n_1, m_1) \cong_{\text{ref}} W(k, n_2, m_2)$ (with $n_i < m_i$ and coprime) implies $(n_1, m_1) = (n_2, m_2)$, it suffices to show that $W_{k, n_1, m_1}^+ \cong W_{k, n_1, m_1}^+$ implies that $(n_1, m_1) = (n_2, m_2)$.

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups

Thank you for your attention !



Torus knot groups,
Garside groups,
complex reflection
groups

Thomas Gobet

Dehornoy-Digne-
Godelle-Krammer-
Michel's monoid
and some torus
knot groups

A new Garside
structure on torus
knot groups and
some braid groups
of complex
reflection groups

"Reflection"
quotients of torus
knot groups